

Bayes Theorem

$$P(h|D) = \frac{P(D|h)P(h)}{P(D)}$$

- $P(h)$ = prior probability of hypothesis h
- $P(D)$ = prior probability of training data D
- $P(h|D)$ = probability of h given D
- $P(D|h)$ = probability of D given h

Choosing Hypotheses

$$P(h|D) = \frac{P(D|h)P(h)}{P(D)}$$

Generally want the most probable hypothesis given the training data

Maximum a posteriori hypothesis h_{MAP} :

$$\begin{aligned} h_{MAP} &= \arg \max_{h \in H} P(h|D) \\ &= \arg \max_{h \in H} \frac{P(D|h)P(h)}{P(D)} \\ &= \arg \max_{h \in H} P(D|h)P(h) \end{aligned}$$

If assume $P(h_i) = P(h_j)$ then can further simplify, and choose the *Maximum likelihood* (ML) hypothesis

$$h_{ML} = \arg \max_{h_i \in H} P(D|h_i)$$

Bayes Theorem

Does patient have cancer or not?

A patient takes a lab test and the result comes back positive. The test returns a correct positive result in only 98% of the cases in which the disease is actually present, and a correct negative result in only 97% of the cases in which the disease is not present. Furthermore, .008 of the entire population have this cancer.

$$\begin{array}{ll} P(cancer) = & P(\neg cancer) = \\ P(+|cancer) = & P(-|cancer) = \\ P(+|\neg cancer) = & P(-|\neg cancer) = \end{array}$$

Basic Formulas for Probabilities

- *Product Rule*: probability $P(A \wedge B)$ of a conjunction of two events A and B:

$$P(A \wedge B) = P(A|B)P(B) = P(B|A)P(A)$$

- *Sum Rule*: probability of a disjunction of two events A and B:

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B)$$

- *Theorem of total probability*: if events A_1, \dots, A_n are mutually exclusive with $\sum_{i=1}^n P(A_i) = 1$, then

$$P(B) = \sum_{i=1}^n P(B|A_i)P(A_i)$$

Brute Force MAP Hypothesis Learner

1. For each hypothesis h in H , calculate the posterior probability

$$P(h|D) = \frac{P(D|h)P(h)}{P(D)}$$

2. Output the hypothesis h_{MAP} with the highest posterior probability

$$h_{MAP} = \operatorname{argmax}_{h \in H} P(h|D)$$